

Top-pions and single top production at HERA and THERA

F. Larios^a, F. Peñuñuri^a, M. A. Pérez^b

^a*Departamento de Física Aplicada, CINVESTAV-Mérida, A.P. 73, 97310 Mérida, Yucatán, México*

^b*Departamento de Física, CINVESTAV, Apdo. Postal 14-740, 07000 México*

We study single top quark production at the HERA and THERA colliders as coming from the FCNC vertices $tc\gamma$ and tcZ that appear at one loop level in Topcolor-assisted technicolor models (TC2). In contrast with previous expectations, we find that the production cross section is of order 10^{-6} pb for HERA and 10^{-4} pb for THERA (even lower than the SM prediction). Therefore, none of the two colliders are capable to probe the pseudoscalar top-pion or the scalar top-higgs predicted in TC2 models via single top production.

PACS numbers: 14.80.-j; 14.65.Ha; 12.60.Nz

1. Introduction

There has been an increased interest in studying forbidden or highly suppressed processes as they become an ideal site to search for new physics lying beyond the Standard Model (SM). In particular, it is expected that if there is any new physics associated with the mass generation mechanism, it may be more apparent in the top quark interactions than in the all the lighter fermions[1]. Along this line, it has been suggested that t-channel single top production could be rather sensitive to the non-SM flavor changing neutral coupling (FCNC) vertices $tc\gamma(Z)$ [2, 3]. While in general, vector and scalar FCNC vertices are very suppressed by the GIM mechanism in the SM, some extensions of the SM can generate some of them at the tree level[4]. This is the case of top-color assisted technicolor (TC2), where a composite $t\bar{t}$ scalar (and a pseudo-scalar) appears as a result of a strong interaction and can give rise to sizeable FCNC $tc\pi_t^0$ ($tc\pi_t^0$) couplings[5]. Then, at the one loop level, these couplings can give rise to effective tcV couplings ($V = \gamma, Z$).

In particular, the pseudo-scalar flavor diagonal and the flavor changing couplings that are relevant for the TC2 prediction of single top quark production are[5, 6]:

$$\begin{aligned} & \frac{m_t(1-\epsilon)}{\sqrt{2}F_t} \frac{\sqrt{v_W^2 - F_t^2}}{v_W} i\bar{t}\gamma^5 t \pi_t^0 \\ & \frac{m_t}{\sqrt{2}F_t} \frac{\sqrt{v_W^2 - F_t^2}}{v_W} i k_{UR}^{tc} k_{UL}^{tt*} \bar{t}_L c_R \pi_t^0, \end{aligned} \quad (1)$$

where $F_t \sim 50\text{GeV}$ is the top-pion decay constant, $v_W = v/\sqrt{2}$, and the k_U terms come from the diagonalization of the up and down quark matrices. Their numerical values are $k_{UL}^{tt} \sim 1$ and $k_{UR}^{tc} \leq \sqrt{2\epsilon - \epsilon^2}$, with $0.03 \leq \epsilon \leq 0.1$ a free parameter. There are also couplings involving the scalar top-higgs h_t^0 , and their effect on the production of single top at HERA is similar to that of the top-pion. In order to simplify our discussion we will not consider them in this work. Notice that in this model the flavor changing scalar coupling only involves the right handed component of the charm quark[6].

Recently, it was suggested that the neutral pseudo-scalar top-pion π_t^0 (or the neutral scalar top-higgs h_t^0) contribution to the single top quark production cross section $\sigma(ec \rightarrow et)$ could be significant, of order 1-6 pb, at the HERA and THERA colliders for a top-pion mass m_π between 200 and 400 GeV[7].

In the present letter, we re-examine this possibility and we find that the cross section due to the top-pion is rather of order much less than 1 fb. In fact, the contribution of the top-pion (or the top-higgs) is much smaller than the SM contribution which is of order less than 1 fb and is not observable at HERA [8, 9]. Unfortunately, none of these two colliders HERA and THERA are thus capable to probe the TC2 scalar particles through single top production.

2. The FCNC single top production at HERA

We consider the tree level FCNC process

$$e(K_1) + c(P_1) \rightarrow e(K_2) + t(P_2) \quad (2)$$

where momentum conservation dictates $K_1 + P_1 = K_2 + P_2$, and the Mandelstam variables are defined as $\hat{s} = (K_1 + P_1)^2$, $t = (P_2 - P_1)^2$ and $u = (P_2 - K_1)^2$. In this work, we take the top quark mass $m_t = 175\text{GeV}$ and the charm quark as

massless. We also neglect terms proportional to the electron mass m_e in the differential cross section. However, we will use $m_e = 0.5\text{MeV}$ for the upper and lower limits of the Mandelstam variable t in the phase space integral. This is done in order to avoid the divergence that appears when the exchanged photon goes on-shell ($t = q^2 = 0$)[9]. The lower and upper limits for $\hat{s} = xS$ are $m_t^2 \leq \hat{s} \leq S$. S is the C.M. energy of the collider: $S = 320\text{GeV}$ for HERA and $S = 1\text{TeV}$ for THERA.

Through a series of loop diagrams the FCNC $t c \pi_t^0$ vertex gives rise to effective $t c V$ vertices, with $V = \gamma, Z$. These diagrams have been calculated and their results given in terms of the following general effective $t c V$ couplings[7]:

$$\Lambda_{V\bar{t}c}^\mu = ie \left(\gamma^\mu (F_{1V} + F_{2V}\gamma^5) - P_2^\mu (F_{3V} + F_{4V}\gamma^5) + P_1^\mu (F_{5V} + F_{6V}\gamma^5) \right). \quad (3)$$

(In the notation of [7] $P_1 = P_c$ and $P_2 = -P_t$.) The F coefficients are given in Ref.[7]. We have computed them and have found agreement, except for the fact that the left handed component of the c quark cannot participate in the $t c \gamma$ vertex. The reason for this is that it is only the right handed component of the c quark that appears in the $t c \pi_t^0$ vertex, which gives rise to $t c \gamma$. Nevertheless, this correction does not change in any way the fact that the cross section for single top production turns out to be negligible for our TC2 model.

3. General $t c \gamma$ vertex with current conservation

For this t -channel process it is well known that HERA will get the bulk of the production cross section from the region where the momentum transfer is very small and the exchanged photon is quasi-real[9]. The contribution from Z exchange is several orders of magnitude smaller. In fact, the experimental situation is such that the scattered positron usually escapes through the rear beam pipe, and events with Q^2 greater than 1GeV^2 are rejected[10]. The reason for this is because the photon propagator tends to infinity in the limit $q^2 \rightarrow 0$, and the phase space integration must be done carefully. In particular, the effective $t c \gamma$ couplings of Eq. (3) are not the most convenient to work with. It turns out that the coefficient F_1 is different from zero in the on-shell limit $t = q^2 = 0$ (see Eq. (5) below). This means that if taken isolated (disregarding the interference with F_3 and F_5) its contribution to the cross section diverges very rapidly. Indeed, the contribution of the other coefficients F_3 and F_5 will also diverge, but in such a way as to cancel out the contribution from F_1 . Unlike the strategy followed by Ref. [7], we will not use the coupling in the form of Eq. (3). Instead, we will perform a Gordon decomposition to transform it into a more suitable form; one that has no such big cancellations.

With the bulk of the contribution coming from a quasi real photon we should bear in mind that electromagnetic vector current conservation implies that the coupling $\bar{t} \gamma_\mu c$ must vanish when the photon goes on shell[11]. In order to make this apparent, we take the $t c \gamma$ coupling of Eq. (3) and change it from the $(\gamma^\mu, P_1^\mu, P_2^\mu)$ basis to the $(\gamma^\mu, \sigma^{\mu\nu} q_\nu, q^\mu)$ basis[11]. We re-write Eq. (3) through a Gordon decomposition:

$$\Lambda_{V\bar{t}c}^\mu = ie \left(\gamma^\mu (V_V - A_V \gamma^5) + i \sigma^{\mu\nu} q_\nu (F_{7V} + F_{8V} \gamma^5) + q^\mu (F_{9V} + F_{10V} \gamma^5) \right) \quad (4)$$

where $q = P_2 - P_1$ is the momentum of the exchange photon (or Z), and with the new coefficients given by

$$\begin{aligned} V &= F_1 + m_t \frac{F_5 - F_3}{2} \\ A &= F_2 + m_t \frac{F_6 - F_4}{2} \\ F_7 &= \frac{F_3 - F_5}{2} \\ F_8 &= \frac{F_4 - F_6}{2} \\ F_9 &= -\frac{F_3 + F_5}{2} \\ F_{10} &= -\frac{F_4 + F_6}{2} \end{aligned}$$

Based on the results of Ref.[7] we can make a numerical calculation of the coefficients defined above. First, let us see the values of F_{3-5} for a top-pion mass $m_\pi = 200\text{GeV}$ and for $q^2 = 0$:

$$\begin{aligned} F_{1\gamma} &= -6.918 \times 10^{-3} f(\epsilon) \\ m_t F_{3\gamma} &= -4.1198 \times 10^{-3} f(\epsilon) \\ m_t F_{5\gamma} &= 9.7162 \times 10^{-3} f(\epsilon). \end{aligned} \quad (5)$$

Here, we have separated the factor $f(\epsilon) = (1 - \epsilon)\sqrt{2\epsilon - \epsilon^2}$ that gives the dependence on ϵ . The cancellation in Eq. (4) can be easily verified. For q^2 small but not zero we have that $V_{\gamma(Z)}$ varies linearly with $t = q^2$. Nevertheless, all the other form factors remain nearly constant for small and even greater (than 1 GeV²) values of t . As mentioned before, the fact that V_γ is directly proportional to q^2 is expected from electromagnetic current conservation[11]. To illustrate this point and to see that the dominant contribution comes from the magnetic transition coefficient $F_{7\gamma}$, let us now note the numerical values of V_γ , $F_{7\gamma}$ and $F_{9\gamma}$ for a top-pion mass $m_\pi = 200\text{GeV}$ and for $-q^2 \leq 1\text{GeV}^2$:

$$\begin{aligned} V_\gamma &= 1.6 \times 10^{-3} \times \frac{t}{m_t^2} \times f(\epsilon) \\ m_t F_{7\gamma} &= -6.9 \times 10^{-3} \times f(\epsilon) \\ m_t F_{9\gamma} &= -2.8 \times 10^{-3} f(\epsilon) \end{aligned} \tag{6}$$

Because V_γ is proportional to $t = q^2$ its contribution to the cross section will not diverge in the $q^2 = 0$ limit. This is not quite the case for F_7 and F_9 ; for them there is still a divergent behaviour, although this time it is only a logarithmic one. On the other hand, the $F_9 \bar{t}q^\mu c$ coupling contribution is proportional to the electron mass, and it is therefore much smaller than that of F_7 .

The coefficients for tcZ are of similar values, even though their contribution is very small we have included them in our numerical results shown in Figs. (1) and (2).

4. Discussion and results

The differential cross section contribution from photon exchange, disregarding terms proportional to the electron mass, is given by[12]:

$$\begin{aligned} \frac{d\sigma}{dt} &= \frac{2\pi\alpha^2}{\hat{s}^2} \frac{M_\gamma}{t^2} \\ M_\gamma &= F_{7\gamma}^2 t [2\hat{s}(m_t^2 - \hat{s}) - m_t^4 - t(2\hat{s} - m_t^2)] \end{aligned} \tag{7}$$

As mentioned in Ref.[12] the bulk of the cross section is given by the photon exchange diagram (M_γ). This is because of its logarithmic divergent behavior that is taken under control with the electron mass. However, our numerical results include the (negligible) contribution from Z exchange.

The total cross section is given by:

$$\sigma = \int_{x_{min}}^1 dx \phi(x) \int_{t_{min}}^{t_{max}} dt \frac{d\sigma}{dt}$$

with $\phi(x)$ the charm quark PDF and $x_{min} = (m_t + m_e)^2/S$ the minimum value of x . The C.M. energy of the collider is $S = 320\text{GeV}$ for HERA and $S = 1\text{TeV}$ for THERA.

The limits for $t = -Q^2$ are:

$$\begin{aligned} t_{min(max)} &= \frac{m_t^4}{4\hat{s}} - (k_{1cm} \pm k_{2cm})^2 \\ k_{1cm} &= \sqrt{\frac{(\hat{s} + m_e^2)^2}{4\hat{s}} - m_e^2} \\ k_{2cm} &= \sqrt{\frac{(\hat{s} + m_e^2 - m_t^2)^2}{4\hat{s}} - m_e^2} \end{aligned} \tag{8}$$

We have evaluated the total cross section with the CTEQ6M PDF[13], running at a fixed scale $\mu = m_t$. We have also run $\phi(x)$ with the energy scale and have seen no significant change.

In Figure (1) we show the production cross section for the process $ec \rightarrow et$ coming from the FCNC vertices $tc\gamma$ and tcZ . For a $m_\pi = 200\text{GeV}$ and $\epsilon = 0.08$ we obtain a tiny cross section $\sigma = 0.5 \times 10^{-6}$ pb, which is even smaller than the SM contribution to single top production. As mentioned before, the dominant contribution comes from the photon diagram so that σ is proportional to $F_{7\gamma}^2$. We can compare with Ref [12], when they take $\kappa_\gamma = 0.1$ for $ec \rightarrow et$ at HERA they obtain a very small cross section of order 0.002pb including NNLO QCD corrections. They instead prefer to discuss $eu \rightarrow et$ because of the small sea charm quark pdf density. In our model, the $t\tau\pi_t^0$ coupling is very small and it will not generate a sizeable $t\tau\gamma$ effective vertex. Also, notice that in our notation we have $m_t F_{7\gamma} = 2\kappa_\gamma$. In




Figure 1: Single top production cross section from π_t^0 as a function of m_π for the HERA collider.

Fig (1) we see that for $\epsilon = 0.08$ and $m_\pi = 200\text{GeV}$ the cross section is $0.5 \times 10^{-6}\text{pb}$. This is four orders of magnitude smaller than the cross section of Ref. [12] because in our case $m_t F_{7\gamma} = 0.0025$ which corresponds to $\kappa_\gamma = 0.001$ two orders of magnitude smaller. This is only a rough comparison, as we do not include NNLO QCD corrections.

In figure (2) we show the production cross section for the THERA collider. It is 4-5 orders of magnitude greater than the one obtained at HERA, but still too small for an experiment with estimated luminosity of 500 pb^{-1} .

As a final remark, let us compare with the results of Ref. [7]. There, the cross section values are 6 orders of magnitude greater. As mentioned before, the cross section in Ref. [7] was calculated from an effective coupling in the form of Eq. (3). This means that a very large cancellation must take place in the small q^2 region, and any standard numerical integration may not be able to handle this situation properly. Rather, the numerical integration is likely to become unstable for small t . Perhaps this is the reason why a (very large) cut in $t_{min}(= -0.001\text{GeV}^2)$ had to be applied in their calculation. In this work, we have instead used the effective coupling in the form of Eq. (4). This new coupling does not induce the large cancellations of the previous one. The numerical integration is stable, even in the low $t = q^2$ region. We use the limits given in Eq. (8), for which t can reach small values of order $t = -10^{-8}\text{ GeV}^2$.

In conclusion, we have shown that neither HERA nor THERA can probe the effects of a top-pion (or top-higgs) of TC2 models via single top quark production.




Figure 2: Single top production cross section from π_t^0 as a function of m_π for the THERA collider.

Acknowledgments

We want to thank C.-P. Yuan for helpful discussions. We thank Conacyt for support.

-
- [1] For a review see D. Chakraborty, J. Konigsberg, D. Rainwater, Annu. Rev. Part. Nucl. Sci. **53** (2003) 301 (hep-ph/0303092); M. Beneke, I. Efthymiopoulos, M. Mangano, J. Womersley et.al., *Top Quark Physics: 1999 CERN Workshop on the SM Physics (and more) at the LHC* (hep-ph/0003033).
 - [2] A. Belyaev, *Single Top quark in the SM and Beyond*, Talk presented at the 8th Intl. Workshop on DIS and QCD (DIS2000), Liverpool, England, April 25-30, 2000. (hep-ph/0007058)
 - [3] T. Han, R.D. Peccei and X. Zhang, Nucl. Phys. B **454** (1995) 527; T. Tait and C.-P. Yuan, Phys. Rev. D **55** (1997) 7300.
 - [4] J.L. Diaz-Cruz, R. Martinez, M.A. Pérez and A. Rosado, Phys. Rev. D **41** (1990) 891; G. Eilam, J.L. Hewett, A. Soni, Phys. Rev. D **44** (1991) 1473.
 - [5] C.T. Hill, Phys. Lett. B **345** (1995) 483; K. Lane, Phys. Lett. B **433** (1998) 96; G. Burdman and D. Kominis, Phys. Lett. B **403** (1997) 101; R.S. Chivukula, E.H. Simmons, J. Terning, Phys. Lett. B **331** (1984) 383; D.J. Muller, S. Nandi, Phys. Lett. B **383** (1996) 345; E. Malkawi, T. Tait and C.-P. Yuan, Phys. Lett. B **385** (1996) 304; R.S. Chivukula, B.A. Dobrescu, H. Georgi and C.T. Hill, Phys. Rev. D **59** (1999) 075003; H. Georgi and A.K. Grant, Phys. Rev. D **63** (2001) 015001.
 - [6] H.-J. He and C.-P. Yuan, Phys. Rev. Lett. **83** (1999) 28; G. Burdman, Phys. Rev. Lett. **83** (1999) 2888.
 - [7] C. Yue, H. Zong and W. Wang, Nucl. Phys. B **667** (2003) 349.
 - [8] U. Baur, Nucl. Phys. B **304** (1988) 451.
 - [9] H. Fritzsch and D. Holtmannspotter, Phys. Lett. B **457** (1999) 186.
 - [10] ZEUS collaboration, Phys. Lett. B **559** (2003) 153; see also, H1 collaboration, Eur. Phys. J. **C33** (2004) 9.

- [11] N.G. Deshpande and M. Nazerimonfared, Nucl. Phys. B **213** (1983) 390.
- [12] N. Kidonakis and A. Belyaev, JHEP 0312 (2003) 004; *ibid.* Phys. Rev. D **65** (2002) 037501.
- [13] CTEQ collaboration. H.L. Lai, et al. JHEP 0207 (2002) 012.